

Implicit Differentiation

1969
AB5

If $3x^2 + 2xy + y^2 = 2$, then the value of $\frac{dy}{dx}$ at $x = 1$ is

- (A) -2 (B) 0 (C) 2 (D) 4 (E) not defined

1969
AB24

If $\sin x = e^y$, $0 < x < \pi$, what is $\frac{dy}{dx}$ in terms of x ?

- (A) $-\tan x$ (B) $-\cot x$ (C) $\cot x$ (D) $\tan x$ (E) $\csc x$

1973
AB40

If $\tan(xy) = x$, then $\frac{dy}{dx} =$

- (A) $\frac{1 - y \tan(xy) \sec(xy)}{x \tan(xy) \sec(xy)}$ (B) $\frac{\sec^2(xy) - y}{x}$ (C) $\cos^2(xy)$
(D) $\frac{\cos^2(xy)}{x}$ (E) $\frac{\cos^2(xy) - y}{x}$

1985
A613

If $x^2 + xy + y^3 = 0$, then, in terms of x and y , $\frac{dy}{dx} =$

- (A) $-\frac{2x+y}{x+3y^2}$ (B) $-\frac{x+3y^2}{2x+y}$ (C) $\frac{-2x}{1+3y^2}$ (D) $\frac{-2x}{x+3y^2}$ (E) $-\frac{2x+y}{x+3y^2-1}$

1988
A69

If $x + 2xy - y^2 = 2$, then at the point $(1, 1)$, $\frac{dy}{dx}$ is

- (A) $\frac{3}{2}$ (B) $\frac{1}{2}$ (C) 0 (D) $-\frac{3}{2}$ (E) nonexistent

1993
AB4

If $x^3 + 3xy + 2y^3 = 17$, then in terms of x and y , $\frac{dy}{dx} =$

(A) $-\frac{x^2 + y}{x + 2y^2}$

(B) $-\frac{x^2 + y}{x + y^2}$

(C) $-\frac{x^2 + y}{x + 2y}$

(D) $-\frac{x^2 + y}{2y^2}$

(E) $\frac{-x^2}{1 + 2y^2}$

1997
AB17

If $x^2 + y^2 = 25$, what is the value of $\frac{d^2y}{dx^2}$ at the point $(4, 3)$?

(A) $-\frac{25}{27}$

(B) $-\frac{7}{27}$

(C) $\frac{7}{27}$

(D) $\frac{3}{4}$

(E) $\frac{25}{27}$

1998
AB6

If $x^2 + xy = 10$, then when $x = 2$, $\frac{dy}{dx} =$

(A) $-\frac{7}{2}$

(B) -2

(C) $\frac{2}{7}$

(D) $\frac{3}{2}$

(E) $\frac{7}{2}$