

1969 AB

13. The region bounded by the x -axis and the part of the graph of $y = \cos x$ between $x = -\frac{\pi}{2}$ and $x = \frac{\pi}{2}$ is separated into two regions by the line $x = k$. If the area of the region for $-\frac{\pi}{2} \leq x \leq k$ is three times the area of the region for $k \leq x \leq \frac{\pi}{2}$, then $k =$

- (A) $\arcsin\left(\frac{1}{4}\right)$ (B) $\arcsin\left(\frac{1}{3}\right)$ (C) $\frac{\pi}{6}$
- (D) $\frac{\pi}{4}$ (E) $\frac{\pi}{3}$

23. The area of the region bounded by the curve $y = e^{2x}$, the x -axis, the y -axis, and the line $x = 2$ is equal to

- (A) $\frac{e^4}{2} - e$ (B) $\frac{e^4}{2} - 1$ (C) $\frac{e^4}{2} - \frac{1}{2}$
- (D) $2e^4 - e$ (E) $2e^4 - 2$

25. A region in the plane is bounded by the graph of $y = \frac{1}{x}$, the x -axis, the line $x = m$, and the line $x = 2m$, $m > 0$. The area of this region

- (A) is independent of m .
- (B) increases as m increases.
- (C) decreases as m increases.
- (D) decreases as m increases when $m < \frac{1}{2}$; increases as m increases when $m > \frac{1}{2}$.
- (E) increases as m increases when $m < \frac{1}{2}$; decreases as m increases when $m > \frac{1}{2}$.

26. $\int_0^1 \sqrt{x^2 - 2x + 1} \, dx$ is

- (A) -1
- (B) $-\frac{1}{2}$
- (C) $\frac{1}{2}$
- (D) 1
- (E) none of the above

27. If $\frac{dy}{dx} = \tan x$, then $y =$

(A) $\frac{1}{2} \tan^2 x + C$

(B) $\sec^2 x + C$

(C) $\ln|\sec x| + C$

(D) $\ln|\cos x| + C$

(E) $\sec x \tan x + C$

29. $\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin x} dx =$

(A) $\ln\sqrt{2}$

(B) $\ln\frac{\pi}{4}$

(C) $\ln\sqrt{3}$

(D) $\ln\frac{\sqrt{3}}{2}$

(E) $\ln e$

33. What is the average (mean) value of $3t^3 - t^2$ over the interval $-1 \leq t \leq 2$?

- (A) $\frac{11}{4}$ (B) $\frac{7}{2}$ (C) 8 (D) $\frac{33}{4}$ (E) 16

35. At $t = 0$ a particle starts at rest and moves along a line in such a way that at time t its acceleration is $24t^2$ feet per second per second. Through how many feet does the particle move during the first 2 seconds?

- (A) 32 (B) 48 (C) 64 (D) 96 (E) 192

38. $\int \frac{x^2}{e^{x^3}} dx =$

(A) $-\frac{1}{3} \ln e^{x^3} + C$

(B) $-\frac{e^{x^3}}{3} + C$

(C) $-\frac{1}{3e^{x^3}} + C$

(D) $\frac{1}{3} \ln e^{x^3} + C$

(E) $\frac{x^3}{3e^{x^3}} + C$

40. If n is a non-negative integer, then $\int_0^1 x^n dx = \int_0^1 (1-x)^n dx$ for

(A) no n

(B) n even, only

(C) n odd, only

(D) nonzero n , only

(E) all n

41. If $\begin{cases} f(x) = 8 - x^2 & \text{for } -2 \leq x \leq 2, \\ f(x) = x^2 & \text{elsewhere,} \end{cases}$ then $\int_{-1}^3 f(x) dx$ is a number between

- (A) 0 and 8 (B) 8 and 16 (C) 16 and 24 (D) 24 and 32 (E) 32 and 40

43. $\int \sin(2x+3) dx =$

- (A) $\frac{1}{2} \cos(2x+3) + C$ (B) $\cos(2x+3) + C$ (C) $-\cos(2x+3) + C$
(D) $-\frac{1}{2} \cos(2x+3) + C$ (E) $-\frac{1}{5} \cos(2x+3) + C$

1969 BC

4. $\int_0^8 \frac{dx}{\sqrt{1+x}} =$

(A) 1

(B) $\frac{3}{2}$

(C) 2

(D) 4

(E) 6

10. $\int_0^1 \frac{x^2}{x^2+1} dx =$

(A) $\frac{4-\pi}{4}$

(B) $\ln 2$

(C) 0

(D) $\frac{1}{2}\ln 2$

(E) $\frac{4+\pi}{4}$

12. If $F(x) = \int_0^x e^{-t^2} dt$, then $F'(x) =$

(A) $2xe^{-x^2}$

(B) $-2xe^{-x^2}$

(C) $\frac{e^{-x^2+1}}{-x^2+1} - e$

(D) $e^{-x^2} - 1$

(E) e^{-x^2}

22. If $f(x) = \int_0^x \frac{1}{\sqrt{t^3+2}} dt$, which of the following is FALSE?

(A) $f(0) = 0$

(B) f is continuous at x for all $x \geq 0$.

(C) $f(1) > 0$

(D) $f'(1) = \frac{1}{\sqrt{3}}$

(E) $f(-1) > 0$